

Treatment Made Simple With XDEMVY®!



Every drop is important! Stay on top of your treatment with XDEMVY.

Start Date: _

One drop in each eye, twice a day (~12 hours apart), for 6 weeks.

Set an alarm and track your doses!



IMPORTANT: Take XDEMVY for the entire 6-week course of treatment.

- If you're using other topical eye medications, ensure a gap of at least 5 minutes between each
- Remove contact lenses and wait 15 minutes after applying XDEMVY before reinserting
- Consult with your doctor about continuing your eyelid hygiene routine

Over time, mites may come back so it's important to see your eye doctor regularly.

	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday	Sunday
	AM	AM	AM	AM	AM	AM	AM
Week 1	PM	PM	PM	PM	PM	PM	PM
	AM	AM	AM	AM	AM	AM	AM
Week 2	PM	PM	PM	PM	PM	PM	PM
	AM	AM	AM	AM	AM	AM	AM
Week 3	PM	PM	PM	PM	PM	PM	PM
	AM	AM	AM	AM	AM	AM	AM
Week 4	PM	PM	PM	PM	PM	PM	PM
	AM	AM	AM	AM	AM	AM	AM
Week 5	PM	PM	PM	PM	PM	PM	PM
	AM	AM	AM	AM	AM	AM	AM
Week 6	PM	PM	PM	PM	PM	PM	PM
Month 9	Follow up with eye doctor						

Indications and Usage

XDEMVY (lotilaner ophthalmic solution) 0.25% is indicated for the treatment of *Demodex* blepharitis.

Important Safety Information

Most common side effects: The most common side effect in clinical trials was stinging and burning in 10% of patients. Other side effects in less than 2% of patients were chalazion/hordeolum and punctate keratitis.

Date & Time:

Handling the Container: Avoid allowing the tip of the dispensing container to contact the eye, surrounding structures, fingers, or any other surface in order to minimize contamination of the solution. Serious damage to the eye and subsequent loss of vision may result from using contaminated solutions.

When to Seek Physician Advice: Immediately seek a physician's advice concerning the continued use of XDEMVY if you develop an intercurrent ocular condition (e.g., trauma or infection), have ocular surgery, or develop any ocular reactions, particularly conjunctivitis and eyelid reactions.

Use with Contact Lenses: XDEMVY contains potassium sorbate, which may discolor soft contact lenses. Contact lenses should be removed prior to instillation of XDEMVY and may be reinserted 15 minutes following its administration.

To report SUSPECTED ADVERSE REACTIONS, contact Tarsus Pharmaceuticals at 1-888-421-4002 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see full <u>Prescribing Information</u>.

By prescription only.

